Assessment for Rectal Diazepam-Diastat Acudial® Medication Administration

Name:

Please circle the correct answer and take the completed test to the school nurse or school district administrator for scoring.

- 1. The three threes refers to counting to three while performing all of the following activities <u>except</u>:
 - a. Before removing syringe from rectum.
 - b. Hold the buttock after withdrawal of the syringe.
 - c. Delivery of the medication in the rectum.
 - d. Insertion of the syringe in the water soluble lubricant.
- 2. If alone with student, what immediate action is recommended after the administration of rectal diazepam?
 - a. Encourage the student to become physically active
 - b. Call emergency medical services or 9-1-1
 - c. Feed the student a snack with carbohydrate and protein foods
 - d. Monitoring the student's gait
- 3. Before administration of rectal diazepam, school personnel should make sure the applicator has the ready collar exposed on the barrel of the syringe.
 - a. True
 - b. False
- 4. After administering the medication, observe the student for the following:
 - a. Walking
 - b. Breathing
 - c. Eating
 - d. You do not need to monitor the student
- 5. When documenting the event, include the following information:
 - a. Student's last meal eaten
 - b. Date, time of seizure and medication, observations
 - c. What the child was wearing
 - d. When the child was last seen at the doctor's office

Number correct: _____ of 5 answers